

Hygromycin B (50 mg/mL)

Product Information

| Product Name | Cat# | Size |
|-------------------------|-----------|----------------|
| Hygromycin B (50 mg/mL) | 60224ES03 | 1 g (20 mL) |
| | 60224ES10 | 10×1 g (20 mL) |

Product Description

Hygromycin B, an aminoglycoside antibiotic synthesized by *Streptomyces hygroscopicus*, inhibits protein synthesis by interfering with 70S ribosomal translocation and inducing misreading of mRNA template, thus killing prokaryotes (such as bacteria), eukaryotes (such as yeasts, fungi) and higher mammalian eukaryotes.


Hygromycin resistance genes (*hyg* or *hph*) derived from *Escherichia coli* encode hygromycin B phosphotransferase, which detoxifies hygromycin B into a non-biologically active phosphorylated product, making it an excellent selective marker for screening and culturing prokaryotic or eukaryotic cells successfully transfected with hygromycin resistance genes. In addition, due to different modes of action, Hygromycin B is often used in combination with G418 or Blasticidin S for selection of dual-resistant cell lines. Hygromycin B can also be used as an antiviral agent because it selectively penetrates into cells with increased membrane permeability due to viral infection and inhibits translation. It can also act as insect-repellant via mixed with animal feed.

This product is sterile hygromycin B solution (50 mg/mL in PBS buffer) and can be directly diluted for use with culture medium.

Product Properties

| | |
|-------------------|--|
| CAS No. | 31282-04-9 |
| Molecular formula | C ₂₀ H ₃₇ N ₃ O ₁₃ |
| Molecular weight | 527.52 g/mol |
| Purity | >90% (HPLC) |
| Potency | ≥1050 U/mg |

Structure



Shipping and Storage

The product is shipped with ice pack and can be stored at 4°C or -20°C for 2 years. Aliquot the solution and avoid repeated freezing and thawing.

Cautions

- Hygromycin B resistance gene (*hyg* or *hph*), except from *E. Coli.*, are also found in other bacterial strains, including *Streptomyces hygroscopicus* and *Klebsiella pneumoniae*.
- Toxic compound, avoid skin and eyes contact, please handle with care.
- For your safety and health, please wear lab coats and disposable gloves for operation.
- For research use only!

Instructions

1. Commonly used screening concentration

The working concentration of hygromycin B for screening stable strains varies according to cell type, culture medium, growth conditions and cell metabolic rate. It is recommended to establish a kill curve (dose-response curve) to determine the optimal screening concentration for the first time. And the recommended working concentration is 50-1000 µg/mL.
Generally, mammalian cells: 50-500 µg/mL; Bacterial/plant cells: 20-200 µg/mL; Fungi: 300-1000 µg/mL.

2. Establishment of killing curve

【Note】 In order to screen stable cell lines, it is necessary to determine the minimum concentration of antibiotics capable of killing untransfected host cells. This can be achieved by establishing a killing curve (dose-response curve). At least five concentrations should be arranged.

1) Day 1: Untransformed cells are planked in an appropriate culture plate at a cell density of 20-25% and cultured overnight.

▲【Note】▲ The amount of inoculation cells can be increased for cells requiring higher density to detect vitality.

2) Set the concentration gradient within the appropriate range according to the cell type. Mammalian cell can set 50, 100, 250, 500, 750, 1000 µg/mL. Dilute the Hygromycin B solution 1:10 with deionized water or PBS buffer to 5 mg/mL. And then dilute the solution to the corresponding working concentration according to the following table.

| Final Concentration (µg/mL) | Medium Volume (mL) | Addition Volume of 5 mg/mL Hygromycin B (mL) |
|-----------------------------|--------------------|--|
| 50 | 9.9 | 0.1 |
| 100 | 9.8 | 0.2 |
| 250 | 9.5 | 0.5 |
| 500 | 9.0 | 1.0 |
| 750 | 8.5 | 1.5 |
| 1000 | 8.0 | 2.0 |

3) Day 2: Replace with a freshly prepared medium containing the corresponding concentration of the drug. Make three parallel samples for each concentration.

4) Then replace with fresh media containing drugs every 3-4 days.

5) Living cell counts are performed at a fixed cycle (e.g., every 2 days) to determine the appropriate concentration. Choose the minimum concentration which kills the majority of cells within an ideal number of days (usually 7-10 days), as the working concentration for screening stable cell line.

3. Screening of stable cell lines

1) 48 h after transfection, the cells were subcultured by screening medium containing hygromycin B at appropriate concentration (direct or diluted).

【Note】 : Antibiotics work best on actively dividing cells. If the cells are too dense, the antibiotic will not kill the cells. Split the cells such that the cells are no more than 25% confluent.

2) Change the screening medium every 3-4 days.

3) Measure the cell colony-formation after 7 days of screening. Colony formation may take another week or more, depending on host cell type, transfection, and screening effectiveness.

4) Pick 5-10 resistant clones and transfer to 35 mm cell culture plates, and cultured with drug-containing screening medium for 7 days.

5) Replace with fresh medium without drugs for culture.

设置格式[10784]: 段落间距段前: 0.5 行, 段后: 0.5 行

设置格式[10784]: 段落间距段前: 0.3 行, 段后: 0.3 行

删除[zuolj2021@outlook.com]:

设置格式[10784]: 字体: 六号

设置格式[10784]: 字体: 六号